



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

Search

PubMed

for

Limits

Preview/Index

History

Clipboard

Details

Preview

Go

Clear

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

- Search History will be lost after eight hours of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search

Most Recent Queries

Time Result

#7 Search exonic enhancers

08:41:27 83

#5 Search exonic silencer

08:40:26 25

#4 Search Intronic silencer

08:39:47 30

#1 Search intronic splicing silencers

08:25:47 3

Clear History

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Jul 17 2003 11:42:11



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search for

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: FEBS Lett. 1998 Oct 16;437(1-2):137-41.

[Related Articles, Links](#)

Regulation of the fibronectin EDA exon alternative splicing. Cooperative role of the exonic enhancer element and the 5' splicing site.

Muro AF, Iaconcig A, Baralle FE.

International Centre for Genetic Engineering and Biotechnology,
Trieste, Italy.

Alternatively spliced exons generally contain weak splicing sites, and exonic and/or intronic regulatory elements recognised by trans-acting auxiliary splicing factors. The EDA exon of the fibronectin gene is a typical example of an exon bearing a purine-rich exon splicing enhancer (ESE) element recognised by members of the SR phosphoprotein family. The regulatory region that governs splicing in the human EDA exon also contains an exon splicing silencer (ESS) element. We have cloned the mouse EDA genomic region, and we show that the ESE and the ESS elements, although they have base differences, can be replaced by the human elements without significant change in the exon inclusion/exclusion ratio. This fact suggests a common splicing regulatory mechanism across species. We demonstrate in vivo the functional activity of the mouse ESE element in splicing. We also show that the trans-acting factors recognising this element cooperate with the 5' splicing site of the EDA exon to facilitate proper exon recognition. Indeed, a strong 5' splicing site overrides the ESE function in exon recognition. However, the presence of a strong 3' splicing site is not sufficient to compensate for the absence of the splicing enhancer. Our data provide in vivo evidence of the interplay between the exonic splicing regulatory elements and the splicing sites, leading finally to subtle regulation of alternative splicing.

PMID: 9804187 [PubMed - indexed for MEDLINE]